

## Research article

# Assessment of the Mobile Mental Health Units' effectiveness in Cyclades islands

Aikaterini Lykomitrou,<sup>1</sup> Stelios Stylianidis,<sup>2</sup> Mary Geitona,<sup>1</sup>  
Stella Pantelidou,<sup>3</sup> Kyriakos Souliotis<sup>1</sup>

<sup>1</sup>Department of Social and Education Policy, University of Peloponnese, Corinth

<sup>2</sup>Department of Psychology, Panteion University of Social and Political Sciences, Athens

<sup>3</sup>Scientific Association for Regional Development and Mental Healthcare (EPAPSY), Greece

**ARTICLE HISTORY:** Received 26 February 2020/Revised 14 July 2020/Published Online 5 August 2021

### ABSTRACT

In Greece, the provision of mental health shows inefficiencies in remote and inaccessible areas due to the lack of appropriate structures and access to healthcare. The purpose of this study was to assess the effectiveness of the Mobile Mental Health Units (MMHUs) in Cyclades with and without MMHUs' operation based on Real-World Evidence (RWE). The study population consisted of 724 people who visited the MMHUs of the western and northeastern Cyclades in 2015. The data derived from the patients' medical records of EPAPSY classified by International Statistical Classification of Diseases and Related Health Problems 10th Revision (ICD-10) and the Global Assessment of Functioning (GAF) scale. The analysis revealed that 60.9% of the participants were women and the average age was 50.1. 50.4% of the people who visited MMHUs without referral from primary health care professionals and 18.8% with referral. The calculation of effectiveness was based on DALYs (Disability-Adjusted Life Years) and was performed according to the World Health Organization methodology. In the specific population, there are no recorded deaths caused by mental disorders and thus DALYs are equal to Years Lost due to Disability (YLDs) with MMHUs' operation. 18% of the population was diagnosed with mood disorders (F30-F39) and morbidity burden 9.49 (YLDs), while 17.5% of the patients were diagnosed with neurotic, stress-related and somatoform disorders (F40-F48) and corresponding morbidity burden 4.53 (YLDs). Our results revealed that the effectiveness of MMHUs corresponds to 17.98 Disability-Adjusted Life Years (DALYs) averted in 2015. The mood disorders and the neurotic, stress-related disorders have shown a high morbidity burden. Advanced age, non-permanent employment, existence of psychopathology in the family and referral on patients' own will were found to significantly affect the mental health status of the participants. The implementation and expansion of flexible and alternative community-based interventions, such as MMHUs, constitute a best practice both for obtaining higher clinical outcomes and for facing regional inefficiencies related to population's access to healthcare.

**KEYWORDS:** Mobile Mental Health Units (MMHUs), Disability-Adjusted Life Years (DALYs), mental health, remote areas, Greece.

### Introduction

Health policy experts and analysts have identified deficiencies in the access and provision of the mental health services in remote and inaccessible areas, in Greece and elsewhere.<sup>1-6</sup> The operation of the community-based<sup>7,8</sup> Mobile Mental Health Units (MMHUs) has been shown to be a cost-effective option for decision-making pur-

poses internationally.<sup>9-22</sup> The Scientific Association for the Regional Development and Mental Health (EPAPSY) launched in 2003 the MMHUs in the Northeastern and Western Cyclades in the islands of Paros, Antiparos, Syros, Tinos, Mykonos, Andros, Kea, Kythnos, Serifos, Sifnos, Kimolos and Milos. These units provide free mental health services to children, adolescents and adults,

are funded by the Ministry of Health and are also supported by the local government and other local agencies.<sup>23,24</sup>

It's worth mentioning that in the above islands and before the MMHUs' operation, patients were facing major difficulties in access to mental health services, with adverse consequences on their health condition.<sup>25</sup> Mental care was only delivered at the outpatient units of the Hospital of Syros. The geographical dispersion among the islands, the difficulties in transportation related with the low frequency of itineraries and the unpredictable weather conditions hampered disproportionately patients' access to health services.<sup>23</sup> In addition, the unpropitious fiscal situation in the country had created a climate of uncertainty, unemployment, financial difficulties, etc., which are directly related to the impact of psychiatric morbidity.<sup>6,26-34</sup>

Throughout these years plenty of actions have taken place by the MMHUs covering the islands' population in need and also providing preventive and consultation mental health services. In particular, workshops and training seminars addressed to doctors, teachers, police officers, civil servants and other local professional groups have been organized. Furthermore, the creation of volunteer groups and the undertaking of actions addressed to specific groups of the population, such as the elderly, victims of domestic violence, parents, children, etc. is a significant part of the operation of MMHUs.<sup>23</sup>

Given the MMHUs' operation and the lack of mental health infrastructure in these islands as well as the geographical features of the country, it was considered critical to evaluate patients' clinical outcomes. Thus, the purpose of this study is to assess the effectiveness of the MMHUs in the Northeastern and Western Cyclades population with symptoms of mental disorders.

## Material and Method

### Study population

The data were extracted from the patients' medical records (Real-World Evidence, RWE) addressed to the MMHUs of the Northeastern and Western Cyclades. The collected data was anonymous and referred to patients' demographic and socio-economic characteristics, the diagnosis according to the International Statistical Classification of Diseases and Related Health Problems 10th Revision (ICD-10),<sup>35</sup> comorbidities, health services use and the overall cases' management. Figure 1 illustrates the sample's stratification for subsequent analysis. The study was undertaken by the University of Peloponnese and has been approved by the Ethics Committee of EPAPSY.

### Clinical data

Medical records were classified in accordance with the diagnostic categories (ICD-10) and the Global Assessment of Functioning scale (GAF).<sup>36</sup> GAF assesses person's health status and the symptoms of mental disorder that affect his/her daily life. The Disability-Adjusted Life Years Index (DALYs) has been used for the outcomes' assessment. DALYs refer to the present value of future life years lost due to premature mortality and morbidity associated with various mental and physical health conditions.<sup>37-42</sup> DALYs for a disease or health condition are calculated as the sum of Years of Life Lost (YLLs) lost due to premature mortality in the population and Years of Lost due to Disability (YLDs). The methodology of the World Health Organization (WHO) was used to calculate the DALYs:<sup>39-41</sup>

$DALYs = YLLs + YLDs$ , where:

$YLLs = \text{Years of Life Lost due to premature death}$

$YLDs = \text{Years Lost due to Disability}$

YLLs were calculated by the formula:<sup>37-42</sup>

$YLLs = N * L$ , where:

$N = \text{number of deaths}$

$L = \text{standard life expectancy at age of death in years}$

YLDs were estimated by the formula (GBD 2010):<sup>40</sup>

$YLDs = P * D$ , where:

$P = \text{number of prevalent cases (prevalence)}$

$D = \text{disability weight for each disease (0-1)}$

Due to the cases of comorbidities, the calculation was formulated as follows:<sup>39</sup>

$YLD_{1+2} = 1 - (1 - YLD_1) * (1 - YLD_2)$


### Statistical analyses

The statistical program IBM SPSS 22.0 was used for data processing (Statistical Package for Social Sciences). The statistical analysis of the results includes descriptive and inferential statistics (linear regression). In the multiple linear regression, DALYs of the mental and physical illnesses were used as dependent variable while the demographic, socio-economic data, health services' use and the cases' overall management were used as independent variables. Logarithmic transformations were performed and the stepwise method was used. Statistical significance levels were set at 0.05.

## Results

### Characteristics of the sample

The study population (table 1) consisted of 724 people, who addressed the MMHUs in 2015. The majority were women, married, secondary and tertiary education graduates and full-time employees. In addition, the



**Figure 1.** Sample Stratification. \*These specific diagnostic categories of diagnoses were selected because they cause greater burden on the population's health and hospitalizations in mental health structures, as evidenced by data from the Hellenic Statistical Authority (2000, 2010–2014). \*\*The GAFscore rating system evaluates the individual's functionality in daily life. The lower the score, the greater the individual's difficulty to cope with everyday life activities. Persons with GAF score 40–31 may not be able to cope with their professional activities. People with GAF score 30–21 need to be cared by their environment. Persons with GAF score between 20–11 are unable to support themselves and need disability allowance and ongoing care.

**Table 1.** Characteristics of the sample.

		N	%
Gender	Male	283	39.1
	Female	441	60.9
	Total	724	100
Marital status	Single	129	18.1
	Married	447	62.9
	Divorced/ Separated	64	9.0
	Widow	52	7.3
	Domestic Partnership	19	2.7
	Total	711	100
Education	No	12	1.7
	Primary school	136	18.8
	Secondary school	70	9.7
	High school	224	30.9
	IEK/College/Vocational training institute	72	9.9
	Technological Educational Institute	62	8.6
	University/Higher Educational Institute	148	20.4
Total	724	100.0	
Occupation	Full-time	327	45.2
	Part-time/seasonal/unemployed	109	15.0
	Household	156	21.5
	Pupil/Student	4	0.6
	Retired	128	17.7
	Total	724	100.0

mean age of the sample was 50.1 years (SD=17.0 years). The 62.9% of participants was married and the median number of children was 2 (1–2 children). 71.0% of the participants lived with their family, while 13.7% lived alone. The majority (91.6%) of the sample were Greeks and 21.4% were lived in Paros, 16.7% in Mykonos and 15.5% in Tinos. 4.3% of participants were uninsured. In addition, 33.7% of people had visited a mental health professional at some point in the past. Specifically, the mean age of the first visit was 44.0 years (SD=17.5 years). The 20.9% reported cases of psychopathology in the family environment. A small number (2.4%) had been hospitalized for mental disorders with an average number of one hospitalization. 28.0% of the participants had serious health problems and 31.4% had been diagnosed with chronic physical illness.

Table 2 presents the reasons for attendance, as well as the referral methods from 2013 to 2017. The reasons for attendance were mainly related to cases of parental counseling, anxiety, depression and relationship or so-

cial problems. It was found that over time, the majority of participants addressed the MMHUs on their own will. It is also worth noting that this method of referral became an increasing trend over time: in 2013 the patients who addressed the units voluntarily approached 61.8% while the percentage reached 82.6% in 2017. As a result, there was a reduction in the other means of referral. In 2015, out of the 365 participants who addressed voluntarily the MMHUs, the majority mentioned as primary reason for their visit the parent counseling (88 patients), anxiety cases (87 patients), and depression cases (66 patients).

### Clinical characteristics

Table 3 presents the patients' diagnosis. The majority was addressed for counseling, mood and neurotic, stress-related and somatoform disorders. Regarding the individualized care plan, 48.9% of participants followed psychological treatments, 21.8% pharmacotherapy and 12.3% a combination of psychological therapies and pharmacotherapy. Specifically, 24.4% followed treatment with antidepressants and 9.4% anxiolytics.

Based on the patients' diagnosis combined with the mortality (YLDs) and morbidity (YLLs) data, DALYs estimation is presented in the table 4 with and without the MMHUs' operation. Thus, 17.98 DALYs were found for the MMHUs' operation and 115.79 in the case of the non MMHUs' operation (table 4).

The results of the multiple linear regression (table 5) were:

- The older the patients, the greater their DALYs.
- Full-time employees had significantly fewer DALYs than the non-permanent work ones.
- Patients who visited MMHUs because of referral had significantly fewer DALYs than those who voluntarily visited MMHUs.
- Patients with a family history of psychopathology had more DALYs than those without family history of psychopathology.

### Discussion

The operation of the MMHUs in the Northeastern and Western Cyclades contributed to the aversion of about 98 DALYs (115.79–17.98) in 2015. This result is due to the prevention of premature mortality and the reduction of the mental health burden. More specifically, before the MMHUs' operation, there were no suitable structures to care for people with symptoms of mental disorders resulting in deficient treatment. It is clear that the MMHUs health services provision cover a significant part of Primary Health Care (PHC) in the field of mental health. The cooperation among PHC and MMHUs health profes-

**Table 2.** Reasons for attendance and referral methods to MMHUs 2013–2017.

	2013		2014		2015		2016		2017	
	N	%	N	%	N	%	N	%	N	%
Reasons for attendance										
Anxiety	75	18.9%	70	17.9%	138	19.1%	72	23.7%	72	18.9%
Depression	76	19.2%	73	18.6%	105	14.5%	52	17.1%	63	16.5%
Psychosis	12	3.0%	12	3.1%	24	3.3%	11	3.6%	9	2.4%
Dementia	16	4.0%	13	3.3%	28	3.9%	14	4.6%	29	7.6%
Drugs	2	0.5%	2	0.5%	6	0.8%	3	1.0%	0	0.0%
Prescription-	45	11.4%	36	9.2%	79	10.9%	30	9.9%	45	11.8%
Problems in relationship-Social-Other	156	39.4%	114	29.1%	98	13.5%	57	18.8%	83	21.8%
Problem of conduct	4	1.0%	4	1.0%	16	2.2%	3	1.0%	5	1.3%
Parents' counseling	10	2.5%	68	17.3%	230	31.8%	62	20.4%	75	19.7%
Total	396	100.0%	392	100.0%	724	100.0%	304	100.0%	381	100.0%
Referral methods										
Without referral/on their own will	275	61.8%	198	50.3%	365	50.4%	199	65.7%	314	82.6%
Primary health care units	35	7.9%	74	18.8%	136	18.8%	36	11.9%	12	3.2%
Private doctor	17	3.8%	32	8.1%	43	5.9%	10	3.3%	3	0.8%
Local municipality	26	5.8%	27	6.9%	53	7.3%	17	5.6%	14	3.7%
Public authorities (police. school etc.)	72	16.2%	55	14.0%	80	11.0%	19	6.3%	14	3.7%
Church	7	1.6%	6	1.5%	10	1.4%	5	1.7%	0	0.0%
Other (family. friends etc.)	13	2.9%	2	0.5%	37	5.1%	17	5.6%	23	6.1%
Total	445	100.0%	394	100.0%	724	100.0%	303	100.0%	380	100.0%

**Table 3.** Sample diagnoses.

Diagnoses according to ICD-10	N	%
Organic, including symptomatic, mental disorders [F00–F09]	39	5.4
Mental and behavioural disorders due to psychoactive substance use [F10–F19]	5	0.7
Schizophrenia, schizotypal and delusional disorders [F20–F29]	20	2.8
Mood [affective] disorders [F30–F39]	130	18
Neurotic, stress-related and somatoform disorders [F40–F48]	127	17.5
Behavioural syndromes associated with physiological disturbances and physical factors [F50–F59]	5	0.7
Disorders of adult personality and behavior [F60–F69]	12	1.7
Mental retardation [F70–F79]	4	0.6
Disorders of psychological development [F80–F89]	1	0.1
Factors influencing health status and contact with health services, counseling [Z03–Z99]	279	38.5
Examination for administrative purposes [Z00–Z02]	62	8.6
Unspecified mental disorder [F99]	40	5.5
Total	724	100.0

sionals would significantly contribute to the continuation of treatment and access to mental care as well as the overall improvement of patients' health status.

Our results indicate that the highest DALYs' scores are found in the diagnostic categories of mood disorders (DALYs: 9.49) and neurotic, stress-related disorders

(DALYs: 4.53) under the MMHUs' operation. These findings are similar to a previous research conducted in Greece indicating high numbers of patients diagnosed with neurotic disorders and mood disorders.<sup>6</sup> It is essential to point out that the results of our study converge with other studies conducted in Greece. In particular,

**Table 4.** DALYs measurement with and without MMHUs' operation.

Diagnostic Categories ICD-10	With MMHUs' operation			Without MMHUs' operation			
	N	YLD	YLL	DALYs	YLD	YLL	DALYs
Organic, including symptomatic, mental disorders [F00–F09]	39	1.17	0.00	1.17	1.69	0.00	1.69
Mental and behavioural disorders due to psychoactive substance use [F10–F19]	5	0.03	0.00	0.03	0.04	0.00	0.04
Schizophrenia, schizotypal and delusional disorders [F20–F29]	20	0.43	0.00	0.43	0.49	79.40	79.89
Mood [affective] disorders [F30–F39]	130	9.49	0.00	9.49	15.60	0.00	15.60
Neurotic, stress-related and somatoform disorders [F40–F48]	127	4.53	0.00	4.53	10.79	0.00	10.79
Behavioural syndromes associated with physiological disturbances and physical factors [F50–F59]	5	0.05	0.00	0.05	0.05	0.00	0.05
Disorders of adult personality and behavior [F60–F69]	12	0.13	0.00	0.13	0.13	0.00	0.13
Mental retardation [F70–F79]	4	0.00	0.00	0.00	0.00	0.00	0.00
Disorders of psychological development [F80–F89]	1	0.00	0.00	0.00	0.00	0.00	0.00
Unspecified mental disorder [F99]	40	0.24	0.00	0.24	0.41	0.00	0.41
Examination for administrative purposes [Z00–Z02]	62	0.15	0.00	0.15	0.65	0.00	0.65
Factors influencing health status and contact with health services, counseling [Z03–Z99]	279	1.75	0.00	1.75	6.53	0.00	6.53
Total	724	17.98	0.00	17.98	36.39	79.40	115.79

**Table 5.** Results of multiple linear regression. Predictors of DALYs.

		$\beta^*$	SE**	P
Age		0.001	0.0001	<0.001
Occupation	Household/Student/Pupil/Retired (refer.)			
	Part-time/Seasonal/Unemployed	–0.004	0.004	0.287
	Full-time/stable	–0.009	0.003	0.003
Referral	No (refer.)			
	Yes	–0.007	0.002	0.006
Existent psychopathology in the family	No (refer.)			
	Yes	0.007	0.003	0.015

\*beta coefficient, \*\*standard error

our finding revealing that patients with psychotic disorders show a higher degree of comorbidity compared to those diagnosed with other mental disorders is also presented in a study conducted in Ioannina-Thesprotia.<sup>43,44</sup> Obviously, these patients have a greater difficulty in finding appropriate treatment and the presence of MMHUs is considered as of added value for the continuation of their treatment. Also, relevant studies have shown that women outnumber men in terms of the onset of mental disorders symptoms,<sup>6,30,31,34</sup> married<sup>30,31,34</sup> as well as people with higher education and higher income have higher incidence of mental disorders.<sup>6</sup> Our results support the above findings since 60.9% of the participants who addressed the MMHUs were women, 62.9% were

married and 20.6% were scientists, businessmen, senior public and private sector executives.

Advanced age, non-permanent employment, existence of psychopathology in the family (heredity) as well patients' referral on their own will, are factors that seem to significantly affect individuals' mental health condition. These findings are in agreement with the international literature that shows increased risk of developing psychopathology and morbidity in these specific population groups.<sup>6,23,26,27,29,31–34,45</sup>

Although DALYs for clinical evaluation of interventions is revealed as among the most valid indices used in the field of mental health internationally,<sup>46–50</sup> it lags far behind

in Greece. The present study measures the effectiveness of an intervention using the DALYs index for the first time in the field of mental health in the country. However, the lack of comparative chronological data on DALYs and the use of only one year for the evaluation of MMHUs constitutes a methodological limitation that should be pointed out. Therefore, DALYs calculation for one year period is indicative and cannot be reproduced over time. The existence of chronological real world evidence data would provide more reliable and robust results.

The results of this study highlighted the contribution of MMHUs to cover unmet mental medical needs. The

expansion of the activities of MMHUs is deemed necessary and beneficial in a country such as Greece where remote areas are inaccessible. Evaluation of health services should be considered as priority in health policy decision making.

## Acknowledgements

It would have been impossible to carry out this research without the availability of data by EPAPSY. Warm thanks to the EPAPSY's Board of Directors and to those who actively supported the conduction of this study.

## References

- Peritogiannis V, Manthopoulou T, Gogou A, Mavreas V. Mental Healthcare Delivery in Rural Greece: A 10-year Account of a Mobile Mental Health Unit. *J Neurosci Rural Pract* 2017, 8:556–561, doi: 10.4103/jnrp.jnrp\_142\_17
- Cheng F, Kirkbride JB, Lennox BR, Perez J, Masson K, Lawrence K et al. Administrative incidence of psychosis assessed in an early intervention service in England: first epidemiological evidence from a diverse, rural and urban setting. *Psychol Med* 2011, 41:949–958, doi: 10.1017/S0033291710002461
- Daig I, Hinz A, Spauschus A, Decker O, Brähler E. Are urban dwellers more depressed and anxious than the rural population? Results of a representative survey. *Psychother Psychosom Med Psychol* 2013, 63:445–454, doi: 10.1055/s-0033-1341469
- Peen J, Shoevers RA, Beekman AT, Dekker J. The current status of urban-rural areas in psychiatric disorders. *Acta Psychiatr Scand* 2010, 121:84–93, doi: 10.1111/j.1600-0447.2009.01438.x
- Riva M, Bamba C, Curtis S, Gauvin L. Collective resources or local social inequalities? Examining the social determinants of mental health in rural areas. *Eur J Public Health* 2011, 21:197–203, doi: 10.1093/eurpub/ckq064
- Skapinakis P, Bellos S, Koupidis S, Grammatikopoulos I, Theodorakis PN, Mavreas V. Prevalence and sociodemographic associations of common mental disorders in a nationally representative sample of the general population of Greece. *BMC Psychiatry* 2013, 13:163, doi: 10.1186/1471-244X-13-163
- Mechanic D. The scientific foundations of community psychiatry. In: Thornicroft G, Szukler G (ed) *Textbook of Community Psychiatry*. Oxford University Press, New York, 2001
- Tansella M, Thornicroft G. *Mental health outcome measures*. Gaskell, London, 2001
- Lykomitrou A, Geitona M, Stylianidis St, Pantelidou St, Souliotis K. Economic Evaluation of Mobile Mental Health Units in Greece: The Case of Cyclades Islands. *Health* 2020, 12:12:891–903, doi: 10.4236/health.2020.127066
- Sowar K, Thurber D, Vanderploeg J, Haldane E. Psychiatric Community Crisis Services for Youth. *Child Adolesc Psychiatr Clin N Am* 2018, 27:479–490, doi: 10.1016/j.chc.2018.03.002
- Yu S, Hill C, Ricks M, Bennet J, Oriol N. The Scope and Impact of Mobile Health Clinics in the United States: A Literature Review. *Int J Equity Health* 2017, 16:178, doi: 10.1186/s12939-017-0671-2
- Lloyd-Evans B, Paterson B, Onyett S, Brown E, Istead H, Gray R et al. National Implementation of a Mental Health Service Model: A Survey of Crisis Resolution Teams in England. *Int J Ment Health Nurs* 2017, 27:214–226, doi: 10.1111/inm.12311
- Mendez-Rubio M, Giannakopoulos P, Gaillard M, Dan L, Budry P. Equipes mobiles de Psychiatrie de l'Âge Avancé. Création de deux équipes mobiles dans les régions Nord et Ouest vaudoises. *Swiss Archives of Neurology, Psychiatry and Psychotherapy* 2012, 163:65–69
- Bonsack C, Holzer L, Stancu I, Baier V, Samitca M, Charbon Y et al. Psychiatric mobile teams for the three ages of live: The Lausanne experience. *Rev Med Suisse* 2008, 4:1960–1969, PMID: 18847128
- Dimopoulou M, Fanti PM, Kiourktsi B, Louvros K, Zennetou I, Kavvadia A et al. Psychosocial intervention in primary health care. The development of the Mobile Mental Health Unit of Corfu/Lefkada. *Medical Chronicles Northwestern Greece* 2011, 8:44–48
- Garoni D, Sarantidis D, Katsadoros K. Utilization of technological resources within the framework of operation of a Mobile Mental Health Unit. *Psychiatriki* 2016, 27:287–295, doi: 10.22365/jpsych.2016.274.287
- Urben S, Baier V, Geiser E, Senent E, Dutoit F, Holzer L. Equipe Mobile pour Adolescents: Une étude pilote évaluant l'efficacité et l'impact des événements de vie aversifs. *Swiss Archives of Neurology, Psychiatry and Psychotherapy* 2013, 164:206–211
- World Health Organization. Mental health: Belgium: switching to community mental health care (Cited 20 January 2020). Available from [https://www.who.int/mental\\_health/evidence/outreach\\_teams\\_belgium/en/](https://www.who.int/mental_health/evidence/outreach_teams_belgium/en/)
- Abrams Z. *New York is launching mobile units*. American Psychological Association 2018, 49:50
- Fernandez C. Innovative mobile clinics serving children and families of Riverside County with limited access to behavioral health services. *American Journal of Psychiatry Residents' Journal* 2017, 11:10–12
- Lloyd-Evans B, Johnson S. Community alternatives to inpatient admissions in psychiatry. *World Psychiatry* 2019, 18:31–32, doi: 10.1002/wps.20587
- Fils-Aimé JR, Grelotti D, Thérosme T, Kaiser B, Raviola G, Alcindor Y et al. A mobile clinic approach to the delivery of community-based mental health services in rural Haiti. *PLoS One* 2018, 13:e0199313, doi: 10.1371/journal.pone.0199313
- Stylianidis S, Pantelidou S, Poullos A, Lavdas M, Lamnidis N. Mobile Mental Health Units on the islands: The Experience of Cyclades. In: Stylianidis S (ed) *Social and Community Psychiatry: Towards a Critical, Patient-Oriented Approach*. Springer International Publishing, Cham, Switzerland, 2016

24. Pantelidou S, Antonopoulou V, Poullos A, Soumaki J, Stylianidis S. Community Child Psychiatry: The Example of Mobile Mental Health Units in the NE and Western Cyclades. In: Stylianidis S (ed) *Social and Community Psychiatry: Towards a Critical, Patient-Oriented Approach*. Springer International Publishing, Cham, Switzerland, 2016
25. Tylee A, Walters P. Underrecognition of anxiety and mood disorders in primary care: why does the problem exist and what can be done? *J Clin Psychiatry* 2007, 68(Suppl 2):27–30, PMID: 17288504
26. Latsou D, Geitona M. The effects of unemployment and economic distress on Depression Symptoms. *Mater Sociomed* 2018, 30:180–184, doi: 10.5455/msm.2018.30.180-184
27. Latsou D, Geitona M. Predictors of physical and mental health among unemployed people in Greece. *Journal of Health and Social Sciences* 2020, 5, 1:115–126, doi: 10.19204/2020/prdc9
28. Giotakos O, Tsouvelas G, Kontaxakis V. Suicide rates and mental health services in Greece. *Psychiatriki* 2012, 23:29–38
29. Laliotis I, Ioannidis J, Stavropoulou C. Total and cause-specific mortality before and after the onset of the Greek economic crisis: an interrupted time-series analysis. *Lancet* 2016, 1: e56–e65, doi: 10.1016/S2468-2667(16)30018-4
30. Madianos M, Economou M, Alexiou T, Stefanis CN. Depression and economic hardship across Greece in 2008 and 2009: two cross-sectional surveys nationwide. *Soc Psychiatry Psychiatr Epidemiol* 2011, 46:943–952, doi: 10.1007/s00127-010-0265-4
31. Economou M, Angelopoulos E, Peppou LE, Souliotis K, Tzavara C, Kontoangelos K et al. Enduring financial crisis in Greece: prevalence and correlates of major depression and suicidality. *Soc Psychiatry Psychiatr Epidemiol* 2016, 51:1015–1024, doi: 10.1007/s00127-016-1238-z
32. Kontaxakis B, Papaslanis T, Havaki-Kontaxaki M, Tsouvelas G, Giotakos O, Papadimitriou G. Suicide in Greece: 2001–2011. *Psychiatriki* 2013, 24:170–174, PMID: 24185083
33. Fountoulakis KN, Koupidis SA, Grammatikopoulos IA, Theodorakis PN. First reliable data suggest a possible increase in suicides in Greece. *BMJ* 2013, 347, f4900, doi: 10.1136/bmj.f4900
34. Economou M, Charitsi M, Peppou LE, Dioti E, Souliotis K. Mental health in Greece during the economic crisis: socioeconomic determinants of depress. *Archives Hellenic Medicine* 2018, 35(Suppl 1):17–26
35. World Health Organization. International Statistical Classification of Diseases and Related Health Problems, 10th Revision (ICD-10). Version for 2016 (Cited 20 January 2020). Available from <http://apps.who.int/classifications/icd10/browse/2016/en#/>
36. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. Fourth Edition, Text Revision (DSM-IV-TR). American Psychiatric Association, Washington DC, 2000
37. Fox Rushby J, Hanson K. Calculating and presenting disability adjusted life years (DALYs) in cost-effectiveness analysis. *Health Policy Plan* 2001, 16:326–331, doi: 10.1093/heapol/16.3.326
38. Driscoll T, Takala J, Steenland K, Corvalan C, Fingerhut M. Review of estimates of the global burden of injury and illness due to occupational exposures. *Am J Ind Med* 2005, 48:491–502, doi: 10.1002/ajim.20194
39. World Health Organization. WHO methods and data sources for global burden of disease estimates 2000–2016. Department of Information, Evidence and Research WHO, Geneva, March 2018 (Cited 20 January 2020). Available from [https://www.who.int/healthinfo/global\\_burden\\_disease/GlobalDALY\\_method\\_2000\\_2016.pdf](https://www.who.int/healthinfo/global_burden_disease/GlobalDALY_method_2000_2016.pdf)
40. World Health Organization. Metrics: Disability-Adjusted Life Year (DALY) (n.d) (Cited 20 January 2020). Available from [http://www.who.int/healthinfo/global\\_burden\\_disease/metrics\\_daly/en/](http://www.who.int/healthinfo/global_burden_disease/metrics_daly/en/)
41. World Health Organization. The Global Burden of Disease concept (n.d) (Cited 20 January 2020). Available from [https://www.who.int/quantifying\\_ehimpacts/publications/en/9241546204chap3.pdf](https://www.who.int/quantifying_ehimpacts/publications/en/9241546204chap3.pdf)
42. Berger M, Bingefors K, Hedblom Ed, Pashos Ch, Torrance G. *Cost of Health Services, Quality and Results*. Dictionary of ISPOR terms. Scientific Supervision: Geitona M, Hatzikou M, Labrelli D, Yfantopoulos I. Papazisis Publications, Athens, 2017
43. Peritogiannis V, Mavreas V. Community mental health teams in Greece: The paradigm of Mobile Mental Health Units. *Archives Hellenic Medicine* 2014, 31:71–76
44. Morden N, Mistler L, Weeks W, Bartels S. Health care for patients with serious mental illness: Family medicine's role. *J Am Board Fam Med* 2009, 22:187–195, doi: 10.3122/jabfm.2009.02.080059
45. Xie S, Karlsson H, Dalman C, Widman L, Rai D, Gardner RM et al. Family History of Mental and Neurological Disorders and Risk of Autism. *JAMA Netw Open* 2019, 2:e190154, doi: 10.1001/jamanetworkopen.2019.0154
46. Gureje O, Chisholm D, Kola L, Lasebikan V, Saxena S. Cost-effectiveness of an essential mental health intervention package in Nigeria. *World Psychiatry* 2007, 6:42–48, PMC1805717
47. Neumann P, Thorat T, Zhong Y, Anderson J, Farquhar M, Salem M et al. A Systematic Review of Cost-Effectiveness Studies Reporting Cost-per-DALY Averted. *PLoS One* 2016, 11:e0168512, doi: 10.1371/journal.pone.0168512
48. Chisholm D (on behalf of WHO-CHOICE). Choosing cost-effective interventions in psychiatry: Results from the CHOICE programme of the World Health Organization. *World Psychiatry* 2005, 4:37–44, PMID: 16633504
49. Knapp M, Wong G. Economics and mental health: the current scenario. *World Psychiatry* 2020, 19:3–14, doi: 10.1002/wps.20692
50. Sagar R, Dandona R, Gururaj G, Dhaliwal RS, Singh A, Ferrari A et al. The burden of mental disorders across the states of India: the Global Burden of Disease Study 1990–2017. *Lancet Psychiatry* 2020, 7:148–161, doi: 10.1016/S2215-0366(19)30475-4

## Ερευνητική εργασία

# Αξιολόγηση της Αποτελεσματικότητας των Κινητών Μονάδων Ψυχικής Υγείας στις Κυκλάδες

Αικατερίνη Λυκομήτρου,<sup>1</sup> Στέλιος Στυλιανίδης,<sup>2</sup> Μαίρη Γείτονα,<sup>1</sup>  
Στέλλα Παντελίδου,<sup>3</sup> Κυριάκος Σουλιώτης<sup>1</sup>

<sup>1</sup>Τμήμα Κοινωνικής και Εκπαιδευτικής Πολιτικής, Πανεπιστήμιο Πελοποννήσου, Κόρινθος

<sup>2</sup>Τμήμα Ψυχολογίας, Πάντειον Πανεπιστήμιο Κοινωνικών και Πολιτικών Επιστημών, Αθήνα

<sup>3</sup>Εταιρεία Περιφερειακής Ανάπτυξης και Ψυχικής Υγείας (ΕΠΑΨΥ), Ελλάδα

**ΙΣΤΟΡΙΚΟ ΑΡΘΡΟΥ:** Παραλήφθηκε 26 Φεβρουαρίου 2020/Αναθεωρήθηκε 14 Ιουλίου 2020/Δημοσιεύθηκε Διαδικτυακά 5 Αυγούστου 2021

### ΠΕΡΙΛΗΨΗ

Η ύπαρξη ελλείψεων σε απαραίτητες και κατάλληλες δομές του συστήματος παροχής υπηρεσιών ψυχικής υγείας σε απομακρυσμένες και δυσπρόσιτες περιοχές έχει διαπιστωθεί και εντείνεται σημαντικά στη νησιωτική χώρα. Αντικείμενο της παρούσας μελέτης είναι η μέτρηση της αποτελεσματικότητας της λειτουργίας των Κινητών Μονάδων Ψυχικής Υγείας (ΚΜΨΥ) βάσει πραγματικών δεδομένων (Real-World Evidence, RWE) για την αντιμετώπιση περιστατικών με συμπτώματα ψυχικών διαταραχών στις Κυκλάδες. Τον πληθυσμό της μελέτης αποτέλεσαν οι 724 εξυπηρετούμενοι που προσήλθαν στις ΚΜΨΥ των Βορειοανατολικών και Δυτικών Κυκλάδων από τα δώδεκα (12) νησιά ευθύνης κατά το έτος 2015. Τα δεδομένα της ανάλυσης προήλθαν από τους ιατρικούς φακέλους των εξυπηρετούμενων της ΕΠΑΨΥ και ταξινομήθηκαν με βάση τη Διεθνή Στατιστική Ταξινόμηση Νόσων και Συναφών Προβλημάτων Υγείας Δέκατη Αναθεώρηση (ICD-10) και την κλίμακα Global Assessment of Functioning (GAF). Από την ανάλυση προέκυψε ότι το 60,9% των εξυπηρετούμενων ήταν γυναίκες και η μέση ηλικία των εξυπηρετούμενων ήταν 50,1 έτη. Το 50,4% των εξυπηρετούμενων απευθύνθηκε για βοήθεια στις ΚΜΨΥ αυτοβούλως και το 18,8% παραπέμφθηκε από τις μονάδες της Πρωτοβάθμιας Φροντίδας Υγείας (ΠΦΥ). Η μέτρηση της αποτελεσματικότητας βασίστηκε στη χρήση του δείκτη των Σταθμισμένων Ετών Ζωής ως προς την Ανικανότητα (Disability-Adjusted Life Years, DALYs), σύμφωνα με τη μεθοδολογία του Παγκόσμιου Οργανισμού Υγείας. Με δεδομένο ότι στον υπό μελέτη πληθυσμό δεν υπάρχει θάνατος από ψυχική διαταραχή, τα DALYs ισούνται με τα Years Lost due to Disability (YLDs). Το 18% του πληθυσμού διαγνώστηκε με διαταραχές της διάθεσης (F30-F39) και φορτίο νοσηρότητας ίσο με 9,49 (YLDs), ενώ το 17,5% των εξυπηρετούμενων με νευρωσικές διαταραχές, οι οποίες σχετίζονται με άγχος και σωματόμορφες διαταραχές (F40-F48) και αντίστοιχο φορτίο νοσηρότητας 4,53 (YLDs). Η αποτελεσματικότητα της λειτουργίας των ΚΜΨΥ καταγράφεται σε 17,98 DALYs που αποτρέπονται κατά το έτος 2015, όπου οι διαταραχές της διάθεσης και οι νευρωσικές σχετιζόμενες με άγχος διαταραχές εμφανίζουν υψηλό φορτίο νοσηρότητας. Η προχωρημένη ηλικία, η μη σταθερή απασχόληση, η ύπαρξη ψυχοπαθολογίας στην οικογένεια και η παραπομπή με πρωτοβουλία του ασθενή βρέθηκε ότι επηρεάζουν σημαντικά την ψυχική υγεία του ατόμου. Η ανάπτυξη ευέλικτων μορφών παρεμβάσεων που θέτουν στο κέντρο τους την κοινότητα, όπως οι ΚΜΨΥ αποτελεί μια άριστη πρακτική με όρους κλινικής αποτελεσματικότητας και παρέχει λύσεις στο πρόβλημα της γεωγραφικής ιδιαιτερότητας και πληθυσμιακής κατανομής στον ελλαδικό χώρο.

**ΛΕΞΕΙΣ ΕΥΡΕΤΗΡΙΟΥ:** Κινητή Μονάδα Ψυχικής Υγείας (ΚΜΨΥ), Σταθμισμένα Έτη Ζωής ως προς την Ανικανότητα (Disability-Adjusted Life Years, DALYs), ψυχική υγεία, δυσπρόσιτες περιοχές, Ελλάδα.

**Συγγραφέας επικοινωνίας:** Αικατερίνη Λυκομήτρου, Τμήμα Κοινωνικής και Εκπαιδευτικής Πολιτικής, Πανεπιστήμιο Πελοποννήσου, Δαμασκηνού & Κολοκοτρώνη, 201 00 Κόρινθος, Διεύθυνση e-mail: k.lykomitrou@gmail.com