

Journal Pre-proof

The effect of COVID-19 pandemic on mental health and quality of life in the Athens area – Greece

Eugenia Triantafillou, Panagiotis Tsellos, Nikos Christodoulou, Chara Tzavara, Irina Mrvoljak-Theodoropoulou, George N. Christodoulou

DOI: <https://doi.org/10.22365/jpsych.2022.095>

To appear in: Psychiatriki Journal

Received date: 16 August 2022

Accepted date: 14 October 2022

Please cite this article as: Eugenia Triantafillou, Panagiotis Tsellos, Nikos Christodoulou, Chara Tzavara, Irina Mrvoljak-Theodoropoulou, George N. Christodoulou, The effect of COVID-19 pandemic on mental health and quality of life in the Athens area – Greece, Psychiatriki (2022), doi: <https://doi.org/10.22365/jpsych.2022.095>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published during the production process, errors, may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

RESEARCH ARTICLE

The effect of COVID-19 pandemic on mental health and quality of life in the Athens area – Greece

Eugenia Triantafyllou, Panagiotis Tsellos, Nikos Christodoulou, Chara Tzavara, Irina Mrvoljak-Theodoropoulou, George N. Christodoulou

Society of Preventive Psychiatry, Greece

ARTICLE HISTORY: Received 16 August 2022 / Revised 10 November 2022 / Published Online 17 November 2022

ABSTRACT

This study examines the associations of the COVID -19 pandemic stressors with mental health and quality of life (QoL) of the general population in different demographic areas of Athens. The random sample of the study consisted of 602 participants, 389 (64,6%) women and 213 (35,4%) men residing in Attica (Greece). It was conducted with telephone interviews during the first wave of the pandemic. The instruments used were: World Health Organization Quality of Life Instrument (WHOQOL-BREF), Depression Anxiety Stress Scale (DASS – 21), Body Vigilance Scale (BVS), Dimensional Obsessive-Compulsive Scale (Category 1: contamination obsessions -decontamination compulsions) (DOCS) and a set of socio-demographic data and questions on stressors related to the COVID-19 outbreak. A range of statistical analyses were used, including descriptive statistics, examination of the questionnaires' internal consistency, calculating Cronbach's alpha coefficient, as well as multiple linear regression analyses with dependent variables the WHOQOL-BREF, DASS-21, DOCS, and BVS scales. According to the results of our study: a) The greater the extent of negative emotions due to the pandemic and the imposed restrictive measures, such as worry and fear, the higher the score of depression, anxiety, stress, obsessive-compulsive symptomatology, body vigilance and the lower the scores of the WHOQOL domains (Physical Health, Psychological Health, Social Relationships, Environment), b) the fear of being contaminated by the virus seems to have intensified depression-anxiety-stress (DASS-21), obsessive-compulsive (DOCS) and hypochondriacal (BVS) symptomatology. c) The independent variables of depression, stress and anxiety and obsessive - compulsive and hypochondriacal symptomatology were negatively associated with QoL d) The most vulnerable groups in terms of QoL and mental health indicators in our study were those with psychiatric or/and physical illness, the elderly, the unemployed during the lockdown period, those with low educational/ socioeconomic status and those living alone. In conclusion, the negative emotions associated with the COVID-19 outbreak and the imposed restrictive measures had a serious impact on mental health and QoL of the population. This and other similar findings should be taken into account by authorities and decision-makers to prevent and deal with the effects of the pandemic. Unfortunately, the COVID-19 pandemic is still continuing and the imposition of new restrictive measures is considered. This increases the relevance of research like the one presented here.

KEYWORDS: COVID-19 stressors, quality of life, depression, stress, contamination obsessions -decontamination compulsions, body vigilance.

Corresponding Author: Eugenia Triantafyllou, Society of Preventive Psychiatry, 52 Vasilissis Sofias Avenue, 115 28, Athens, Greece, Email: etrianta@med.uoa.gr

Introduction

The sudden coronavirus pandemic has spread at an alarming rate worldwide provoking millions of deaths. Especially in Greece, a total of 4336 cases had been confirmed as of July 30th 2020 and the number of deaths was 203 whereas on November 29th 2020 the number of confirmed cases was 104.227 and the number of accumulated deaths was 2.321. Up to now (July 2022), over 30.000 people passed away due to COVID-19.

Beyond the physical impacts, the COVID-19 pandemic had various serious mental health effects. Indeed, relevant studies and meta-analyses have shown high levels of psychological distress in the general population, with higher levels of anxiety and depression in people with underlying diseases, and in those actually infected by the virus,¹ as well as post-traumatic stress symptoms.² The rapid spread of the disease, its high mortality rate and concerns about the future are probably the main sources of anxiety.³ In addition, according to the findings of several studies, the fear and distress linked to the virus spread has contributed to high levels of depression, anxiety,^{4,5,6} obsessive compulsive disorder (OCD) symptomatology as well as health anxiety.⁷ Moreover, the concern about COVID 19 was positively correlated with both obsessive compulsive symptomatology (OCS) and health anxiety symptoms.⁸

Pandemics overall have a significant impact on individuals' quality of life (QoL) in different dimensions such as psychological and physical health, social and family relationships, issues related to work-unemployment etc.^{9,10,11} Indeed, research referring to the impact of coronavirus disease and related restrictions on mental health and QoL of the general population in Austria, showed increase of depression and decrease of QoL in times of COVID-19 as compared to the pre- COVID-19 period.¹²

Also, as mentioned in a recent review,¹³ those who have been quarantined, have reported a significantly high prevalence of psychological distress, emotional disturbance, depression, stress, low mood, irritability, insomnia, posttraumatic stress, anger and emotional exhaustion. It seems that the lockdown, the loss of normal routine and reduced social and physical contact with others are stressors that probably contribute to boredom, irritability and isolation from the rest of the world.

The present study was conducted by the Society of Preventive Psychiatry. We investigated the impact of the COVID-19 outbreak and of the related imposed restrictive measures on mental health and QoL of the general population in Attica, Greece. The objectives of the study were:

- To assess QoL, stress-anxiety-depression, contamination obsessions/ decontamination compulsions and the degree of body vigilance.
- To investigate the associations of QoL with stress-anxiety-depression, contamination obsessions/ decontamination compulsions and the degree of body vigilance.
- To detect the effect of sociodemographic characteristics and of stressors related to the COVID-19 outbreak on QoL and on the rest of the variables of the study.

Materials and methods

Participants and procedures

The sample consisted of 602 persons, 389 (64.6%) women and 213 (35.4%) men with mean age 47.8 (SD= 17.0) years. Table 1 provides descriptive summaries of the demographic characteristics of the sample. Our research was carried out during the first period of restrictions imposed in Greece to minimize the spread of the pandemic. It was carried out using telephone interviews, with the application of systematic sampling. Specifically, a

random sample was taken from several districts in Athens, by selecting 5 citizens' telephone numbers every 10 pages of the Attica region telephone directory, maintaining the anonymity of personal data. The interviews were conducted during a three-month time span (from April to June 2020). The number of 602 participants refers to the individuals who answered our phone calls and accepted to participate in our research during the above time span.

Table 1

Measures

The instruments used were:

*World Health Organization QOL Instrument -WHOQOL-BREF:*¹⁴ The Greek version of WHOQOL – BREF includes 30 questions assessing an overall quality of life/general health facet, and 4 domains: physical health, psychological health, social relationships and environment. This instrument has been adapted in Greek population¹⁵ with satisfactory internal consistency (Cronbach's alpha coefficient: 0,67-0,81) and satisfactory construct, convergent and discriminant validity. In our study, calculation of Cronbach's alpha coefficient per domain was performed, which resulted in satisfactory alpha values ranging from 0.70–0.76.

*Depression Anxiety Stress Scale- DASS – 21:*¹⁶ It includes 21 questions. The depression, scale measures: dysphoria, hopelessness, devaluation of life, self-deprecation, lack of interest/involvement, anhedonia and inertia. The anxiety scale estimates autonomic arousal, skeletal musculature effects, situational anxiety and subjective experience of anxious affect. The stress scale measures: difficulty relaxing, persistent arousal, agitation, irritability/overreactiveness and impatience. The DASS was shown to possess satisfactory psychometric properties, and the factor structure was substantiated both by exploratory and confirmatory factor analysis.¹⁶ This instrument has been adapted in Greek population¹⁷ with satisfactory internal consistency for the total scale (Cronbach's alpha coefficient: 0,93) and satisfactory psychometric properties concerning validity. In our study, calculation of Cronbach's alpha coefficient per each subscale was performed, which resulted in satisfactory alpha values ranging from 0.83–0.93.

*Body Vigilance Scale –BVS:*¹⁸ a four-item questionnaire measuring conscious attention focused on internal bodily sensations. There are three items that assess the degree of Attentional focus on bodily sensations, the Perceived sensitivity to changes in bodily sensations, and the Average amount of time spent attending to bodily sensations. The fourth item involves separate ratings for Attention to 15 sensations (e.g., heart palpitations). An overall score is also calculated.

The Body Vigilance Scale has been validated in clinical and nonclinical samples with satisfactory psychometric properties (reliability-validity).¹⁸ BVS scale underwent a rigorous translation in Greek, back translation and cross-examination by bilingual subjects. In our study, calculation of Cronbach's alpha coefficient for BVS total score was performed, which resulted in satisfactory alpha value (0.75).

*Dimensional Obsessive-Compulsive Scale – DOCS:*¹⁹ assessing contamination obsessions and decontamination compulsions (Category 1: Concerns about Germs and Contamination), which includes 5 questions and an overall score, acquiring satisfactory psychometric properties (reliability and validity).¹⁹ DOCS (Category 1) underwent a rigorous translation in Greek, back translation and cross-examination by bilingual subjects. In our study, calculation of Cronbach's alpha coefficient for DOCS and especially for the subscale "Concerns about Germs and Contamination" was performed, which resulted in satisfactory alpha value (0.80). *Set of socio-demographic data and questions-stressors related to the COVID-19 outbreak*, which refer to: sex, age, residence, educational status, years of education, professional status, income, family status, number of children, living circumstances, socioeconomic status, job status during quarantine, job condition, category of respondents regarding COVID-19

infection.

The questions-stressors related to the COVID-19 outbreak evaluate the following: frequency and reasons of movements outside the home, accuracy and frequency of getting mass media information about the virus, negative feelings due to the pandemic and restrictive measures, fear of contamination, negative feelings due to social isolation, insecurity due to job and economic matters.

Statistical analysis

Statistical analyses were conducted using SPSS statistical software (version 22.0). A range of statistical analyses were used, including descriptive statistics and examination of the questionnaires' internal consistency, calculating Cronbach's alpha coefficient. Also, multiple linear regression analyses in a stepwise method (p for entry 0.05, p for removal 0.10) were used with dependent variables the WHOQOL-BREF, DASS-21, DOCS, and BVS scales. The regression equation included also terms for demographic characteristics and stressors regarding the pandemic. The adjusted regression coefficients (β) with standard errors (SE) were computed from the results of the linear regression analyses. When DOCS or BVS scales were the dependent variables, multiple linear regression was conducted after having logarithmically transformed the dependent variable. All reported p values are two-tailed. Statistical significance was set at $p < 0.05$.

Results

Sample characteristics

Regarding the stressful conditions that have been developed due to the COVID-19 and the related restrictive measures: the majority of the participants ($n=497$, 82.6 %) reported that they were satisfied with the accuracy of mass media information about COVID-19 and many of them ($n=238$, 39.5%) were getting information 2-3 times per day. Many participants used to get out of the house during the quarantine period several times per week ($n=202$, 33.6%), getting out mostly for shopping ($n=480$, 79.9%).

The most prevalent negative feelings reported by the majority of the participants were worry ($n=491$, 81.6%), stress ($n=386$, 64.1%) and fear ($n=288$, 47.8%). A significant percentage of the participants ($n=171$, 28.7%) reported high levels of negative feelings due to the social isolation and moderate levels of negative feelings ($n=207$, 34.4%) due to the pandemic and the restrictive measures in general. Also, a significant proportion of the population ($n=222$, 36.9%) exhibited moderate levels of fear of being contaminated and high levels of fear of loved ones getting the virus ($n=232$, 38.5%). Finally, a significant number of participants ($n=260$, 43.2%) reported high/very high levels of insecurity due to financial matters and a restricted proportion of the sample showed high/very high levels of job insecurity ($n=191$, 31.8%).

Associations of WHOQOL- BREF

According to multiple linear regression analysis, men, older people, those with lower socioeconomic status, unemployed participants during the quarantine and those with mental or/and physical illness showed lower scores in the majority of WHOQOL-BREF domains. Also, married people exhibited higher QoL scores.

Concerning the stressors related to the COVID-19 outbreak: the greater the extent of negative emotions due to the pandemic and the restrictive measures, the lower the scores of all QoL domains. Higher scores of the fear of significant others contracting the virus and of occupational insecurity were associated with lower scores in certain QoL domains.

Depression and stress scores were significantly negatively associated with WHOQOL-BREF domain scores. Higher score of average amount of time spent attending to bodily sensations was associated with lower scores in WHOQOL-BREF domains (Table 2)

Associations of DASS-21

According to multiple linear regression analysis, participants with lower score of educational and socioeconomic status and those with mental or/and physical illness exhibited higher scores in DASS-21 domains. Also, people who shared accommodation showed a lower score in significant DASS-21 dimensions.

The greater the extent of negative emotions due to the pandemic and the restrictive measures and the greater the fear of being infected by the virus, the higher the scores of DASS-21 dimensions. Participants who reported that they had accurate information about COVID -19 from the media showed lower score in DASS-21 dimensions (Table 3).

Associations of DOCS

According to multiple linear regression analysis, people with lower educational level, those with higher monthly income and patients with mental or/and physical illness, in comparison to healthy individuals, exhibited higher score in DOCS scale. The more frequent updates provided to the participants about COVID-19, the greater the fear of getting the virus and the greater the extent of negative emotions, the higher the score in DOCS scale (Table 4).

Associations of BVS

According to multiple linear regression analysis, older participants, those with higher educational and socioeconomic status, patients with physical or/and mental illness showed higher scores in significant BVS dimensions, in comparison to healthy individuals.

Negative emotions, insecurity for financial matters and the fear of being contaminated by the virus, are positively correlated with BVS scores (Table 5).

Discussion

The present study evaluated the impact of the COVID-19 outbreak and the imposed measures on QoL and mental health of the general population in the Region of Attica, in Greece. In particular, it examined its effect with reference to QoL, stress-anxiety and depression symptomatology as well as contamination obsessions/ decontamination compulsions, and degree of body vigilance.

Men, older people, singles, those with a lower socioeconomic status and participants with physical or/ and mental illness exhibited lower levels in certain QoL domains. These findings are in line with the results of a research in China demonstrating that health related quality of life in the general population deteriorated significantly with aging, with chronic disease and lower income.²⁰ This can be explained by the fact that the elderly are affected disproportionately by the pandemic because they constitute a high risk group for developing severe illnesses.²¹ Also it seems that elderly people have a poor QoL associated with restrictive measures.²² With reference to the differential effect of the pandemic on men and women, contrary to our findings, an Italian research showed that women had an overall worse psychological, physical, and environmental QoL during the pandemic compared to men.²³ Our findings are also consistent with the results of other studies in Saudi Arabia and Morocco, according to which people with chronic medical conditions reported significantly lower QoL scores during COVID-19 pandemic.^{24,25} Accordingly, another study concerning U.S. young adults during the initial months of the COVID-19 pandemic, showed a deteriorated health-related QoL among those with either a suspected or reported mental health diagnosis.²⁶

As expected, the wider the extent of negative emotions arising from the pandemic and the restrictive measures, the more likely was the occurrence of a low level of QoL. Also, the higher the levels of depression, stress and of body vigilance, the more likely it was that the QoL would decline. In accordance with the results of our study, in a cross-sectional survey in

Saudi Arabia, it was demonstrated that participants who had experienced anxiety, depression and stress were at an increased risk of exhibiting lower levels of QoL.²⁴

With reference to the association of depression-anxiety-stress with sociodemographic characteristics, those who resided alone and those with a lower educational and socioeconomic status seemed to exhibit higher levels of stress. Moreover, it was observed that the lower the educational level of the participants, the higher the possibility to have depressive symptomatology. Our findings are in line with the results of some other studies in Greece^{27,28} and in Mexico,²⁹ according to which, persons with lower educational level were more likely to develop depressive and anxiety symptomatology during the pandemic period. Also, participants who suffered from physical or/and psychiatric disorder were more likely to exhibit higher levels of anxiety and stress. These findings are in accordance with the results of an Italian study³⁰ which indicate that persons with a medical history were more likely to have higher scores of anxiety and depression.

According to our findings, the higher the negative feelings due to COVID- 19 and the higher the fear of contamination, the higher the possibility of developing depression, anxiety and stress. This finding is in line with other Greek studies which indicated that young people who worried more of being contaminated by COVID-19, were more likely to develop psychological distress.³¹ Similarly, according to other studies, a perceived high risk of contracting the virus as well as the fear and anxiety about COVID-19 are associated with high levels of depression and stress.^{32,5,6} Our participants who had precise information by the media about COVID-19 had a lower score of depression and stress. This finding is in line with other studies, according to which higher satisfaction with the health information concerning COVID-19 was associated with a lower psychological impact of the outbreak.^{33,29}

In reference to contamination obsession and decontamination compulsions, our findings indicate that unhealthy participants and those with higher income were more likely to develop OCS. In the same direction, according to other studies persons with mental disorders³⁴ and higher income groups were more likely to exhibit higher levels of OCS during the pandemic situation.³⁵ It appears that rich people are probably more “careful” and value their lives more than the less financially privileged.

Also, the more negative emotions (such as fear) the participants developed due to the pandemic, the more likely they were to develop OCS. This is consistent with previous reports maintaining that obsessions and compulsions (as washing rituals) were often motivated by fear of contracting a disease.^{36,37}

Regarding body vigilance, our findings indicated that older participants, those with higher educational status and those suffering from mental or/and physical illness were more likely to develop hypochondriacal symptomatology. In accordance with these findings, other study results indicated that persons with past or present psychiatric illnesses had greater sensitivity to and awareness of their body sensations.³⁸

The great extent of negative emotions developed, the fear of contamination by the virus and the high degree of financial insecurity seems to have intensified hypochondriacal symptomatology. These findings are consistent with the results of other studies^{39,7} that revealed a positive relationship between excessive concern about COVID-19 and health anxiety.

In conclusion, the more negative feelings that the participants developed due to the COVID- 19 pandemic, the higher the possibility of developing depression, anxiety, stress, obsessive- compulsive and hypochondriacal symptomatology. These conditions in turn seem to contributing to the deterioration of QoL.

The main limitation of this study is that the cross-sectional design used limits drawing conclusions regarding causality, therefore further longitudinal studies could capture the ongoing psychological impact of the COVID-19 pandemic and the related effect on QoL over a longer duration of time. Such a longitudinal study is scheduled to be published soon. Also,

another limitation is that the random population, specifically targeted on the Attica region excluding the rest of Greece, does not constitute a national representation.

A positive aspect of our work is that it included the study of the impact of the pandemic not only with reference to depression, anxiety and stress (which has been the subject of most studies), but also with reference to obsessive-compulsive and hypochondriacal symptomatology, as well as to QoL. Quality of life is a broad multi-dimensional indicator and “can be used in daily clinical practice for the assessment of therapeutic interventions and for the health need assessment of populations”^{40,41}, especially in response to disasters and crises such as the COVID-19 pandemic. The findings of our study have demonstrated that the greater the extent of negative emotions due to the pandemic the lower the QoL scores. Also, our study shows the significant effect of negative emotions due to the pandemic and fear of contamination on hypochondriacal and obsessive-compulsive symptomatology, a very important finding since only few researchers^{7,8} have addressed this issue.

Moreover, the sample of our study included not only healthy subjects but also patients with physical and / or mental diseases, thus rendering possible the comparison between these groups. Finally, another positive feature of our study was that it was carried out with telephone interviews, (rather than internet questionnaires), each interview lasting almost one hour and thus allowing clarification of concepts and issues related to the questionnaires.

The findings of this and other related studies should be taken into consideration by the authorities so that appropriate interventions to improve quality of life can be designed and actions to prevent and treat psychopathological conditions associated with the COVID-19 pandemic can be implemented.

The recent Athens Mental Health Summit Declaration of the WHO European Region (July, 2021) “on actions required to address the impact of the COVID-19 pandemic on mental health” is a good response to this need. However, this is just a starting point. It should be followed by substantial attention to the needs and adequate and consistent funding. Finally, vulnerable population groups suffering from mental or/and physical diseases will certainly require specific and targeted support to cope with the negative impact of disasters such as COVID-19 pandemic on their mental health and quality of life.

Acknowledgements

The authors express their appreciation to all the participants volunteering to this study.

References

1. Luo Y, Chua CR, Xiong Z, Ho RC, Ho CS. A systematic review of the impact of viral respiratory epidemics on mental health: an implication on the coronavirus disease 2019 pandemic. *Fron. Psychiatry* 2020, 11:565098, doi.org/10.3389/fpsyt.2020.565098
2. Salari N, Hosseini-Far A, Jalali R, Vaisi-Raygani A, Rasoulpoor S, Mohammadi M et al. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Glob Health* 2020, 16: 1-11, doi: 10.1186/s12992-020-00589-w
3. Banerjee D. The COVID-19 outbreak: Crucial role the psychiatrists can play. *Asian J Psychiatr* 2020, 50: 102014, doi: 10.1016/j.ajp.2020.102014
4. Ahorsu DK, Imani V, Lin CY, Timpka T, Broström A, Updegraff JA et al. Associations between fear of COVID-19, mental health, and preventive behaviours across pregnant women and husbands: an actor-partner interdependence modelling. *Int J Ment Health Addict* 2020, 11: 1-15, doi:10.1007/s11469-020-00340-x
5. Lee SA, Mathis AA, Jobe MC, Pappalardo EA. Clinically significant fear and anxiety of COVID-19: A psychometric examination of the Coronavirus Anxiety Scale. *Psychiatry research* 2020, 290: 113112, doi: 10.1016/j.psychres.2020.113112

6. Lee SA, Crunk EA. Fear and psychopathology during the COVID-19 crisis: neuroticism, hypochondriasis, reassurance-seeking, and coronaphobia as fear factors. *Omega: J. Death Dying* 2020, 85:483-496, doi: 10.1177/0030222820949350
7. Wheaton MG, Prikhidko A, Messner G. Is fear of COVID-19 contagious? The effects of emotion contagion and social media use on anxiety in response to the coronavirus pandemic. *Front. Psychol* 2021, 11:567379, doi: 10.3389/fpsyg.2020.567379
8. Wheaton MG, Messner GR, Marks JB. Intolerance of uncertainty as a factor linking obsessive-compulsive symptoms, health anxiety and concerns about the spread of the novel coronavirus (COVID-19) in the United States. *J of Obsessive-Compuls Relat Disord* 2021, 28:100605, doi: 10.1016/j.jocrd.2020.100605
9. Chen J, Wu J, Hao S, Yang M, Lu X, Chen X et al. Long term outcomes in survivors of epidemic Influenza A (H7N9) virus infection. *Sci Rep* 2017, 7: 17275, doi: 10.1038/s41598-017-17497-6
10. Batawi S, Tarazan N, Al-Raddadi R, Al Qasim E, Sindi A, Al Johni S et al. Quality of life reported by survivors after hospitalization for Middle East respiratory syndrome (MERS). *Health Qual Life Outcomes* 2019, 17: 1-7, doi: 10.1186/s12955-019-1165-2
11. Kwek SK, Chew WM, Ong KC, Ng AW, Lee LS, Kaw G et al. Quality of life and psychological status in survivors of severe acute respiratory syndrome at 3 months post discharge. *J Psychosom Res* 2006, 60: 513-519, doi: 10.1016/j.jpsychores.2005.08.020
12. Pieh C, O' Rourke T, Budimir S, Probst T. Relationship quality and mental health during COVID-19 lockdown. *PLoS One* 2020, 15: e023890, doi: 10.1371/journal.pone.0238906
13. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020, 395.10227: 912–920, doi: 10.1016/S0140-6736(20)30460-8
14. The WHOQOL Group. The World Health Organization Quality of Life Assessment (WHOQOL): Development and general psychometric properties. *Soc Sci Med* 1998, 46: 1569-1585, doi: 10.1016/s0277-9536(98)00009-4
15. Ginieri-Coccosis M, Triantafillou E, Tomaras V, Soldatos C, Mavreas V, Christodoulou. Psychometric properties of WHOQOL-BREF in clinical and healthy Greek populations: Incorporating new culture-relevant items. *Psychiatriki* 2012, 23:130-142. PMID: 22796911
16. Lovibond P, Lovibond SH. The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behav Res Ther* 1995, 33: 335-343, doi: 10.1016/0005-7967(94)00075-u
17. Lyrakos GN, Arvaniti C, Kostopanagiotou G. Depression, Anxiety, Stress Scale 21 (DASS-21). In: Stalikas A, Triliva S, Roussi P (eds) *Psychometric tools in Greece*. Pedio Press, Athens, 2012.
18. Schmidt NB, Lerew DR, Trakowski JH. Body vigilance in panic disorder: Evaluating attention to bodily perturbations. *J Consult Clin Psychol* 1997, 65: 214, doi:10.1037/0022-006X.65.2.214
19. Abramowitz JS, Deacon, BJ, Olatunji BO, Wheaton MG, Berman NC, Losardo D et al. Assessment of obsessive-compulsive symptom dimensions: development and evaluation of the Dimensional Obsessive-Compulsive Scale. *Psychol Assess* 2010, 22: 180-198, doi: 10.1037/a0018260
20. Ping W, Zheng J, Niu X, Guo C, Zhang J, Yang H, Shi Y. Evaluation of health-related quality of life using EQ-5D in China during the COVID-19 pandemic. *PloS one* 2020, 15: e0234850, doi: 10.1371/journal.pone.0234850
21. Mueller AL, McNamara MS, Sinclair DA. Why does COVID-19 disproportionately affect older people? *Aging (albany NY)* 2020, 12: 9959, doi: 10.18632/aging.103344
22. Ferreira LN, Pereira LN, da FéBrás M, Ilchuk K. Quality of life under the COVID-19 quarantine. *Qual Life Res* 2021, 30: 1389-1405, doi: 10.1007/s11136-020-02724-x

23. Epifanio MS, Andrei F, Mancini, G, Agostini F, Piombo MA, Spicuzza V et al. The impact of COVID-19 pandemic and lockdown measures on quality of life among Italian general population. *J. Clin Med* 2021, 10: 289, doi: 10.3390/jcm10020289
24. Algahtani FD, Hassan SU, Alsaif B, Zrieq R. Assessment of the quality of life during COVID-19 pandemic: A cross-sectional survey from the Kingdom of Saudi Arabia. *Int J Environ Res Public Health* 2021, 18: 847, doi: 10.3390/ijerph18030847
25. Samlani Z, Lemfadli Y, Errami AA, Oubaha S, Krati K. The impact of the COVID-19 pandemic on quality of life and well-being in Morocco. *Arch Public Health* 2020, 6 : 130-134, doi: 10.17352/2455-5479.000091
26. Liu CH, Zhang E, Wong GTF, Hyun S. Factors associated with depression, anxiety, and PTSD symptomatology during the COVID-19 pandemic: Clinical implications for US young adult mental health. *Psychiatry Res* 2020, 290: 113172, doi: 10.1016/j.psychres.2020.113172
27. Karaivazoglou K, Konstantopoulou G, Kalogeropoulou M, Iliou T, Vorvolakos T, Assimakopoulos K et al. Psychological distress in the Greek general population during the first COVID-19 lockdown. *BJ Psych Open* 2021, 7: e59, doi: 10.1192/bjo.2021.17
28. Parlapani E, Holeva V, Voitsidis P, Blekas A, Gliatas I, Porfyri GN et al. Psychological and behavioral responses to the COVID-19 pandemic in Greece. *Fron. Psychiatry* 2020, 11 :821, doi: 10.3389/fpsy.2020.00821
29. Cortés-Álvarez NY, Pineiro-Lamas R, Vuelas-Olmos CR. Psychological effects and associated factors of COVID-19 in a Mexican sample. *Disaster Med Public Health Prep* 2020, 14: 413-424, doi: 10.1017/dmp.2020.215
30. Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C et al. A nationwide survey of psychological distress among Italian people during the COVID-19 pandemic: immediate psychological responses and associated factors. *Int J Environ Res Public Health* 2020, 17: 3165, doi:10.3390/ijerph17093165
31. Kornilaki EN. The psychological effect of COVID-19 quarantine on Greek young adults: Risk factors and the protective role of daily routine and altruism. *Int J Psychol* 2022, 57: 33-42, doi: 10.1002/ijop.12767
32. Newby JM, O'Moore K, Tang S, Christensen H, Faasse K. Acute mental health responses during the COVID-19 pandemic in Australia. *PloS one* 2020, 15: e0236562, doi: 10.1371/journal.pone.0236562
33. Wang C, Pan R, Wan X, Tan Y, Xu L, McIntyre RS et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun* 2020, 87: 40-48, doi: 10.1016/j.bbi.2020.04.028
34. Zheng Y, Xiao L, Xie Y, Wang H, Wang G. Prevalence and Characteristics of Obsessive-Compulsive Disorder Among Urban Residents in Wuhan During the Stage of Regular Control of Coronavirus Disease-19 Epidemic. *Front Psychiatry* 2020, 11:1435, doi: 10.3389/fpsy.2020.594167
35. AlHusseini N, Sajid M, Altayeb A, Alyousof S, Alsheikh H, Alqahtani A et al. Depression and Obsessive-Compulsive Disorders Amid the COVID-19 Pandemic in Saudi Arabia. *Cureus* 2021, 13: e12978, doi: 10.7759/cureus.12978
36. Rachman S. Fear of contamination. *Behav Res Ther* 2004, 42: 1227-1255, doi: 10.1016/j.brat.2003.10.009
37. Wheaton MG, Abramowitz JS, Berman NC, Riemann BC, Hale LR. The relationship between obsessive beliefs and symptom dimensions in obsessive-compulsive disorder. *Behav Res Ther* 2010, 48: 949-954, doi: 10.1016/j.brat.2010.05.027
38. Özdin S, BayrakÖzdin Ş. Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: The importance of gender. *Int Soc Psychiatry* 2020, 66: 504–511, doi: 10.1177/0020764020927051

39. Asmundson GJ, Taylor S. How health anxiety influences responses to viral outbreaks like COVID-19: What all decision-makers, health authorities, and health care professionals need to know. *J Anxiety Disord* 2020, 71: 102211. doi: 10.1016/j.janxdis.2020.102211
40. Katschnig H. How useful is the concept of quality of life in psychiatry. In: Katschnig H. Freeman H, Sartorius N (eds) *Quality of Life in Mental Disorders*. John Wiley & Sons, New York, 1999
41. Triantafillou E, Economou M, Christodoulou GN. Quality of life and mental health: Conceptual references and theoretical models. In: Kontaxakis VP, Chavaki-Kontaxaki MI, Christodoulou GN (eds) *Preventive Psychiatry and Mental Hygiene*. Beta Med Publ, Athens, Greece, 2005

Table1. Socio-demographic characteristics

Variables		N	%
Gender	Women	389	64.6
	Men	213	35.4
Age, mean (SD)		47.8 (17.0)	
Number of children	0	316	52.5
	1	108	17.9
	≥2	178	29.5
Number of habitants at home, mean (SD) median (IQR)		1.8 (1.2)	2 (1 - 3)
Years of Education, mean (SD)		15 (3.9)	
Family status	Married	242	40.2
	Living together	50	8.3
	Unmarried	225	37.4
	Widowed	35	5.8
	Divorced	43	7.1
	Separated	7	1.2
Living circumstances	With own family	281	46.7
	With parental family	110	18.3
	Alone	147	24.4
	With partner (unmarried)	64	10.6
Professional status	Full-time employed	315	52.3
	Part-time employed	33	5.5
	Student (graduate/postgraduate)	23	3.8
	Retired	119	19.8
	Household	30	5.0
	Unemployed	41	6.8
	Other	41	6.8
Individuals under suspension of work	No	254	73.0
	Yes	94	27.0
Job condition	At work	150	55.4

Socioeconomic status	Remotely	121	44.6
	Upper class	32	5,3
	Upper middle class	123	20.4
	Middle class	333	55,3
	Middle-lower class	92	15,3
	Lower class	22	3,7

Table 2. Multiple linear regression analysis with WHOQOL- BREF domains as dependent variables

		β +	SE++	b‡	P
<i>Dependent: WHOQOL-BREF Overall QoL/ and general health</i>					
Gender	Women (reference)				
	Men	-2.73	1.17	-0.08	0.020
Age		-0.14	0.05	-0.13	0.006
Socioeconomic status		-2.04	0.79	-0.10	0.010
Group	Healthy (reference)				
	Patients with psychiatric illness	-8.70	2.22	-0.14	<0.001
	Patients with physical illness	-8.23	1.49	-0.19	<0.001
	Patients with psychiatric and physical illness	-8.43	2.64	-0.11	0.002
Negative feelings due to pandemic/restrictive measures		-2.73	0.61	-0.17	<0.001
Depression		-1.10	0.23	-0.25	<0.001
Average amount of time spent attending to bodily sensations		-1.01	0.31	-0.15	0.001
<i>Dependent: WHOQOL-BREF Physical health</i>					
Socioeconomic status		-1.66	0.61	-0.09	0.006
Group	Healthy (reference)				
	Patients with psychiatric illness	-5.58	1.71	-0.10	0.001
	Patients with physical illness	-6.59	1.15	-0.18	<0.001
	Patients with psychiatric and physical illness	-10.89	2.04	-0.17	<0.001
Fear of beloved ones being contaminated		-1.38	0.56	0.09	0.014
Negative feelings due to pandemic/restrictive measures		-1.37	0.47	-0.10	0.004
Depression		-1.51	0.18	-0.39	<0.001
Stress		-0.37	0.16	-0.11	0.025
Average amount of time spent attending to bodily sensations		-1.03	0.24	-0.17	<0.001
<i>Dependent: WHOQOL-BREF Psychological health</i>					
Married/ Living together	No (reference)				
	Yes	2.90	1.13	0.10	0.011
Socio-economic status		-1.26	0.63	-0.07	0.045

Negative feelings due to pandemic/restrictive measures		-1.26	0.49	-0.09	0.010
Depression		-1.79	0.18	-0.48	<0.001
Stress		-0.36	0.17	-0.11	0.037

Dependent: WHOQOL-BREF Social relationships

Age		-0.10	0.05	-0.11	0.046
Married/ Living together	No (reference)				
	Yes	5.38	1.42	0.17	<0.001
Negative feelings due to pandemic/restrictive measures		-1.49	0.61	-0.10	0.015
Depression		-1.41	0.23	-0.35	<0.001

Dependent: WHOQOL-BREF Environment

Socioeconomic status		-2.74	0.61	-0.20	<0.001
Working during quarantine	No (reference)				
	Yes	-2.93	1.21	-0.12	0.016
Negative feelings due to pandemic/restrictive measures		-1.86	0.47	-0.17	<0.001
Insecurity due to job matters		-1.19	0.43	-0.13	0.006
Depression		-0.70	0.18	-0.23	<0.001

+regression coefficient ++Standard Error ‡Standard Regression Coefficient
Note: only significant variables are presented

Table 3. Multiple linear regression analysis with DASS-21 dimensions dependent variables and sociodemographic characteristics and stressors related to COVID-19 as predictors

		β +	SE++	b‡	P
<i>Dependent: DASS21 Depression</i>					
Educational status		-0.20	0.09	-0.10	0.033
Group	Healthy (reference)				
	Patients with psychiatric illness	2.13	0.54	0.15	<0.001
	Patients with physical illness	0.76	0.37	0.08	0.041
	Patients with psychiatric and physical illness	3.55	0.64	0.21	<0.001
Fear of being contaminated		0.38	0.17	0.10	0.030
Negative feelings due to pandemic/restrictive measures		1.10	0.14	0.30	<0.001
<i>Dependent: DASS21 Anxiety</i>					
Educational status		-0.30	0.09	-0.15	0.001
Living circumstances	Alone (reference)				
	With others	-0.76	0.35	-0.09	0.032
Group	Healthy (reference)				

	Patients with psychiatric illness	2.34	0.50	0.18	<0.001
	Patients with physical illness	0.59	0.34	0.07	0.083
	Patients with psychiatric and physical illness	3.33	0.59	0.21	<0.001
Fear of being contaminated		0.63	0.16	0.18	<0.001
Fear of beloved ones being contaminated		0.35	0.17	0.10	0.035
Negative feelings due to pandemic/restrictive measures		0.70	0.13	0.20	<0.001
<i>Dependent: DASS21 Stress</i>					
Educational status		-0.31	0.11	-0.13	0.004
Socioeconomic status		-0.46	0.22	-0.09	0.040
Group	Healthy (reference)				
	Patients with psychiatric illness	2.16	0.61	0.13	<0.001
	Patients with physical illness	0.68	0.42	0.06	0.106
	Patients with psychiatric and physical illness	3.18	0.72	0.16	<0.001
Accurate information about COVID-19	No (reference)				
	Yes	-1.17	0.42	-0.10	0.006
Fear of being contaminated		0.49	0.20	0.12	0.012
Fear of beloved ones being contaminated		0.52	0.20	0.12	0.011
Negative feelings due to pandemic/restrictive measures		1.31	0.16	0.31	<0.001
<i>Dependent: Total DASS-21</i>					
Educational status		-0.82	0.25	-0.14	0.001
Living circumstances	Alone (reference)				
	With others	-2.00	1.01	-0.08	0.049
Socio-economic status		-1.09	0.52	-0.08	0.036
Group	Healthy (reference)				
	Patients with psychiatric illness	6.67	1.43	0.17	<0.001
	Patients with physical illness	2.02	0.98	0.08	0.039
	Patients with psychiatric and physical illness	10.04	1.69	0.21	<0.001
Accurate information about COVID-19	No (reference)				
	Yes	-2.07	0.98	-0.07	0.035
Fear of being contaminated		1.51	0.46	0.15	0.001
Fear of beloved ones being contaminated		1.19	0.48	0.11	0.013
Negative feelings due to pandemic/restrictive measures		3.10	0.37	0.31	<0.001

+regression coefficient ++Standard Error ‡standardized regression coefficient

Note: only significant variables are presented

Table 4. Regression analysis with the total DOCS score as dependent variable and sociodemographic characteristics and stressors related to COVID-19 as predictors

		β+	SE++	b‡	P
Educational status		-0.24	0.08	-0.11	0.002
Monthly income		0.22	0.11	0.07	0.042
Group	Healthy (reference)				
	Patients with psychiatric illness	2.37	0.49	0.17	<0.001
	Patients with physical illness	0.84	0.34	0.09	0.014

	Patients with psychiatric and physical illness	2.13	0.59	0.12	<0.001
Frequency of getting information		0.66	0.17	0.14	<0.001
Fear of being contaminated		1.40	0.13	0.37	<0.001
Negative feelings due to pandemic/restrictive measures		0.65	0.13	0.17	<0.001

+regression coefficient ++Standard Error ‡standardized regression coefficient

Note: only significant variables are presented

Table 5. Multiple regression analysis with BVS as dependent variable and sociodemographic characteristics and stressors related to COVID-19 as predictors

		β +	SE++	b‡	P
<i>Dependent: BVS Attentional focus on bodily sensations</i>					
Age		-0.02	0.01	-0.16	<0.001
Group	Healthy (reference)				
	Patients with psychiatric illness	0.96	0.35	0.11	0.006
	Patients with physical illness	0.84	0.26	0.14	0.001
	Patients with psychiatric and physical illness	1.76	0.44	0.16	<0.001
Fear of being contaminated		0.65	0.09	0.28	<0.001
Insecurity due to economic matters		0.16	0.08	0.07	0.048
<i>Dependent: BVS Perceived sensitivity to changes in bodily sensations</i>					
Age		-0.03	0.01	-0.18	<0.001
Socioeconomic status		-0.35	0.12	-0.11	0.003
Group	Healthy (reference)				
	Patients with psychiatric illness	0.91	0.35	0.10	0.009
	Patients with physical illness	0.81	0.26	0.13	0.002
	Patients with psychiatric and physical illness	2.20	0.44	0.20	<0.001
Fear of being contaminated		0.77	0.09	0.32	<0.001

<i>Dependent: BVS Average amount of time spent attending to bodily sensations</i>					
Age		-0.01	0.01	-0.08	0.050
Group	Healthy (reference)				
	Patients with psychiatric illness	0.76	0.33	0.09	0.025
	Patients with physical illness	0.97	0.25	0.16	<0.001
	Patients with psychiatric and physical illness	1.71	0.43	0.16	<0.001
Fear of being contaminated		0.80	0.09	0.34	<0.001
<i>Dependent: BVS Attention to 15 sensations</i>					
Educational status		0.10	0.04	0.09	0.028
Group	Healthy (reference)				
	Patients with psychiatric illness	1.20	0.28	0.17	<0.001
	Patients with physical illness	0.57	0.20	0.11	0.004
	Patients with psychiatric and physical illness	0.07	0.35	0.12	0.002
Fear of being contaminated		0.30	0.09	0.16	0.001
Fear of beloved ones being contaminated		0.25	0.10	0.12	0.009
Negative feelings due to pandemic/restrictive measures		0.40	0.08	0.21	<0.001
<i>Dependent: BVS Total score</i>					
Age		-0.08	0.02	-0.17	<0.001
Group	Healthy (reference)				
	Patients with psychiatric illness	3.91	1.06	0.14	<0.001
	Patients with physical illness	3.27	0.80	0.16	<0.001

	Patients with psychiatric and physical illness	7.00	1.36	0.20	<0.001
Fear of being contaminated		2.83	0.29	0.37	<0.001

+regression coefficient ++Standard Error ‡standardized regression coefficient

Note: only significant variables are presented

Journal Pre-proof

ΕΡΕΥΝΗΤΙΚΗ ΕΡΓΑΣΙΑ

Η επίδραση της πανδημίας COVID-19 στην ψυχική υγεία και την ποιότητα ζωής του γενικού πληθυσμού στην Αττική

Ευγενία Τριανταφύλλου, Παναγιώτης Τσέλλος, Νίκος Χριστοδούλου, Χαρά Τζαβάρα, Ιρίνα Mrvoljak- Θεοδωροπούλου, Γεώργιος Ν. Χριστοδούλου

Εταιρεία Προληπτικής Ψυχιατρικής

ΙΣΤΟΡΙΚΟ ΑΡΘΡΟΥ: Παραλήφθηκε 16 Αυγούστου 2022/ Αναθεωρήθηκε 10 Νοεμβρίου 2022 / Δημοσιεύθηκε Διαδικτυακά 17 Νοεμβρίου 2022

ΠΕΡΙΛΗΨΗ

Με την παρούσα μελέτη επιδιώκεται η εκτίμηση της επίδρασης στρεσογόνων παραγόντων που σχετίζονται με την πανδημία COVID-19 στην ψυχική υγεία και την ποιότητα ζωής (ΠΖ) του γενικού πληθυσμού σε περιοχές της Αττικής. Το δείγμα της μελέτης, που επελέγη με τυχαία δειγματοληψία, αποτέλεσαν 602 άτομα του γενικού πληθυσμού, κάτοικοι διαφορετικών περιοχών της Αττικής, εκ των οποίων 389 (64,6%) ήταν γυναίκες και 213 (35,4%) άνδρες. Η έρευνα πραγματοποιήθηκε με τηλεφωνικές συνεντεύξεις κατά το πρώτο κύμα της πανδημίας. Τα ερωτηματολόγια που χρησιμοποιήθηκαν ήταν: το Ερωτηματολόγιο Ποιότητας Ζωής του Παγκόσμιου Οργανισμού Υγείας (WHOQOL-BREF), το Ελληνικό Ερωτηματολόγιο Μέτρησης της Κατάθλιψης, του Άγχους και του Στρες (DASS-21), η Κλίμακα εστίασης της προσοχής σε σωματικές αισθήσεις-συμπτώματα (BVS), η Πολυδιάστατη Κλίμακα Ιδεοψυχαναγκαστικής συμπτωματολογίας (Κατηγορία 1: ιδεοληψίες μόλυνσης και καταναγκασμοί απολύμανσης) (DOCS) και ένα δελτίο κοινωνικοδημογραφικών χαρακτηριστικών και ερωτήσεων που αφορούν σε στρεσογόνους παράγοντες σχετιζόμενους με την πανδημία της COVID-19. Σύμφωνα με τα αποτελέσματα της μελέτης: α) Όσο μεγαλύτερη είναι η έκταση των αρνητικών συναισθημάτων εξαιτίας της πανδημίας και των περιοριστικών μέτρων, που επεβλήθησαν για την μείωση της εξάπλωσης της πανδημίας, τόσο υψηλότερα είναι τα επίπεδα της κατάθλιψης, του άγχους, του στρες, της ιδεοψυχαναγκαστικής συμπτωματολογίας, της εστίασης στα σωματικά συμπτώματα και τόσο χαμηλότερο είναι το επίπεδο της ΠΖ, β) Ο φόβος μόλυνσης από τον ιό φαίνεται να επιδεινώνει την κατάθλιψη, το άγχος και το στρες (DASS-21), την ιδεοψυχαναγκαστική (DOCS) και την υποχονδριακή (BVS) συμπτωματολογία, γ) Οι ανεξάρτητες μεταβλητές της κατάθλιψης, του στρες, του άγχους, της ιδεοψυχαναγκαστικής και της υποχονδριακής συμπτωματολογίας φαίνεται να συσχετίστηκαν αρνητικά και να επέδρασαν στην ποιότητα ζωής του πληθυσμού, δ) Οι πιο ευάλωτες ομάδες όσον αφορά την ΠΖ και την ψυχική υγεία ήταν τα άτομα που έπασχαν από ψυχική ή/και σωματική νόσο, οι ηλικιωμένοι, οι άνεργοι κατά την περίοδο του lockdown, τα άτομα χαμηλού εκπαιδευτικού επιπέδου και χαμηλής κοινωνικοοικονομικής κατάστασης και όσοι ζούσαν μόνοι τους. Εν κατακλείδι, τα αρνητικά συναισθήματα που σχετίζονται με την πανδημία της COVID-19 και τα με τα περιοριστικά μέτρα είχαν πιθανώς σοβαρό αρνητικό αντίκτυπο στην ψυχική υγεία και στην ΠΖ του πληθυσμού. Το εύρημα αυτό καθώς και άλλα παρόμοια ερευνητικά πορίσματα θα πρέπει

να αξιοποιηθούν από τις αρχές και τους υπεύθυνους λήψης αποφάσεων, για την πρόληψη και την αντιμετώπιση των επιπτώσεων της πανδημίας. Μιας πανδημίας που δυστυχώς συνεχίζεται και δημιουργεί ήδη προβληματισμό ως προς την ανάγκη εκ νέου εφαρμογής αυστηρών περιοριστικών μέτρων. Κάτω από αυτό το πρίσμα, εργασίες όπως η παρούσα μπορεί να αποδειχθούν χρήσιμες.

ΛΕΞΕΙΣ ΕΥΡΕΤΗΡΙΟΥ: Στρεσογόνοι παράγοντες COVID-19, ποιότητα ζωής, κατάθλιψη, στρες, ιδεοληψίες μόλυνσης και καταναγκασμοί απολύμανσης, εστίαση στα σωματικά συμπτώματα.

Επιμελητής συγγραφέας: Ευγενία Τριανταφύλλου, Εταιρεία Προληπτικής Ψυχιατρικής, Βασιλίσσης Σοφίας 52, 115 281, Αθήνα, Ελλάδα, Email: etrianta@med.uoa.gr

Journal Pre-proof